

Sadílek D.: Štěnice „na paprice“ aneb Jak se vaří obsah jaderné DNA
(Živa 2019, 6: 320–322)

Citovaná a další doporučená literatura:

- Bennett MD, Leitch IJ, Price HJ & Johnston JS 2003: Comparisons with *Caenorhabditis* (~100 Mb) and *Drosophila* (~175 Mb) using flow cytometry show genome size in *Arabidopsis* to be ~157 Mb and thus ~25% larger than the *Arabidopsis* genome initiative estimate of ~125 Mb. *Annals of Botany* **91**: 547–557. doi: 10.1093/aob/mcg057
- Doležel J & Geilhuber J 2010: Nuclear genome size: Are we getting closer? *Cytometry Part A* **77A**: 635–642. doi: 10.1002/cyto.a.20915
- Gregory TR 2019. Animal genome size. Database. <http://www.genomesize.com>
- Hanrahan SJ & Johnston JS 2011: New genome size estimates of 134 species of arthropods. *Chromosome Research* **19**: 809–823. doi: 10.1007/s10577-011-9231-6
- Henry TJ 2017: Biodiversity of Heteroptera, pp. 223–263. In: Footit RG & Adler PH (Eds). *Insect Biodiversity: Science and Society, Volume 1, Second Edition*. Wiley-Blackwell, United Kingdom, pp 987.
- Johnston JS, Ross LD, Beani L, Hughes DP & Kathirithamby J 2004: Tiny genomes and endoreduplication in Strepsiptera. *Insect Molecular Biology* **13**: 581–585. doi: 10.1111/j.0962-1075.2004.00514.x
- Johnston JS, Yoon KS, Strycharz JP, Pittendrigh BR & Clark JM 2007: Body lice and head lice (Anoplura: Pediculidae) have the smallest genomes of any hemimetabolous insect reported to date. *Journal of Medical Entomology* **44**: 1009–1012. doi: 10.1093/jmedent/44.6.1009
- Panzera F, Ferrandis I, Ramsey J, Salazar-Schettino PM, Cabrera M, Monroy C, Bargues MD, Mas-Coma S, O'Connor JE, Angulo VM, et al. 2007: Genome size determination in Chagas disease transmitting bugs (Hemiptera-Triatominae) by flow cytometry. *The American Journal of Tropical Medicine and Hygiene* **76**: 516–521. doi: 10.4269/ajtmh.2007.76.516
- Panzera F, Pérez R, Panzera Y, Ferrandis I, Ferreiro MJ & Calleros L 2010: Cytogenetics and genome evolution in the subfamily Triatominae (Hemiptera, Reduviidae). *Cytogenetic and Genome Research* **128**: 77–87. doi: 10.1159/000298824
- Rodrigues ASB, Silva SS, Pina-Martins F, Loureiro J, Castro M, Gharbi K, Johnson KP, Dietrich CH, Borges PAV, Quartau JA, et al. 2016: Assessing genotype–phenotype associations in three dorsal colour morphs in the meadow spittlebug *Philaneus spumarius* (L.) (Hemiptera: Aphrophoridae) using genomic and transcriptomic resources. *BMC Genetics* **17**: 144. doi: 10.1186/s12863-016-0455-5
- Sadílek D, Šťáhlavský F, Vilímová J & Zima J 2013: Extensive fragmentation of the X chromosome in the bed bug *Cimex lectularius* Linnaeus, 1758 (Heteroptera,

Cimicidae): A survey across Europe. *Comparative Cytogenetics* **7**: 253–269.

doi: 10.3897/CompCytogen.v7i4.6012

Sadílek D, Urfus T, Hadrava J, Vilímová J & Suda J 2019: Nuclear genome size in contrast to chromosome number variability in human bed bug, *Cimex lectularius* (Heteroptera: Cimicidae). *Cytometry Part A* **95A**: 746–756. doi: 10.1002/cyto.a.23729

Suda J & Leitch IJ 2010: The quest for suitable reference standards in genome size research. *Cytometry Part A* **77A**: 717–720. doi: 10.1002/cyto.a.20907