

Šenfěldr M., Maděra P. Hřížení smrku v ekotonu horní hranice lesa (*Živa* 2022, 4: 179–182)

Použitá a doporučená literatura

- Arno, S.F., Hammerly, R.P., 1984: Timberline. Mountain and Arctic Forest Frontiers. The Mountaineers, Seattle, 304 pp.
- Čermák J, Deml M, Penka M (1973) A new method of sap flow rate determination in trees. *Biologia Plantarum* 15:171–178.
- Holtmeier, F.K. 2009. Mountain Timberlines. Ecology, Patchiness, and Dynamics. *Advances in global change research* 36. Springer science. 437 p.
- Jeník J. et Lokvenc, T. 1962. Die alpine Waldgrenze im Krkonoše Gebirge. *Rozpr. Čs.Akad. věd, Praha, ser. Math. - natur.*, 72/1 : 1-65.
- Jeník, J. 1959. Příspěvek k poznání horní hranice rozšíření dřevin ve Vysokých Sudetech. *Acta dendrologica Československa* 1:21 – 30.
- Jeník, J. 1976. O vegetativním rozmnožování smrku, *Picea abies* (L.) Karsten (On vegetative reproduction in spruce, *Picea abies* (L.) Karsten). *BEN* (At F.[ed.]: Štúdie o ihličnatých drevinách (Studies in coniferous trees), Veda, Bratislava, 101-108.
- Jeník, J., 1961. Alpinská vegetace Krkonoš, Kralického Sněžníku a Hrubého Jeseníku. Nakladatelství Československé akademie věd, Praha, 412 s.
- Körner, Ch. 2012. Alpine treelines: Functional ecology of the global high elevation tree limits, Springer, Basel.
- Kučera J, Urban J (2012) History of the development of the trunk heat balance method in last forty years. *Acta Horticulturae* 951:87–94.
- Kučera, J., Urban, J. 2012. History of the development of the trunk heat balance method in last forty years. *Acta Hort.* 951:87-94.
- Larcher, W. 2003. *Physiological plant ecology: ecophysiology and stress physiology of functional groups*. Springer.
- Novák J., Petr L., Treml V. 2010. Late-Holocene human-induced changes to the extent of alpine areas in the East Sudetes, Central Europe. *The Holocene* 20(6) 895 - 905.
- Plesník P. 1971. Horná hranice lesa vo Vysokých a v Belanských Tatrách. Nakl. SAV, 475 s., Bratislava.
- Šenfěldr, M. 2014. Populace smrku ztepilého při horní hranici lesa v Hrubém Jeseníku. Lesnická a dřevařská fakulta Mendelovy univerzity, Dizertační práce, 139 s.
- Šenfěldr, M., & Maděra, P. (2011). Population structure and reproductive strategy of Norway spruce (*Picea abies* L. Karst) above the former pastoral timberline in the Hrubý Jeseník Mountains, Czech Republic. *Mountain Research and Development*, 31(2), 131-143.
- Šenfěldr, M., Kaczka, R., Buras, A., Samusevich, A., Herrmann, C., Spyt, B., ... & Treml, V. (2021). Diverging growth performance of co-occurring trees (*Picea abies*) and shrubs (*Pinus mugo*) at the treeline ecotone of Central European mountain ranges. *Agricultural and Forest Meteorology*, 308, 108608.
- Šenfěldr, M., Treml, V., Maděra, P., & Volařík, D. (2014). Effects of prostrate dwarf pine on Norway spruce clonal groups in the treeline ecotone of the Hrubý Jeseník Mountains, Czech Republic. *Arctic, Antarctic, and Alpine Research*, 46(2), 430-440.

- Šenfeldr, M., Urban, J., Maděra, P., & Kučera, J. (2016). Redistribution of water via layering branches between connected parent and daughter trees in Norway spruce clonal groups. *Trees*, 30(1), 5-17.
- Tranquillini W (1979) Physiological Ecology of the Alpine Timberline. Tree Existence at High Altitudes with Special Reference to the European Alps. Ecological studies 31. Berlin, Germany: Springer.
- Treml, V. 2007. Dynamika alpinské hranice lesa ve Vysokých Sudetech. Doktorská práce, Katedra fyzické geografie a geoekologie, Přírodovědecká fakulta Univerzity Karlovy, Dizertační práce 191 s.
- Treml, V., & Banaš, M. (2008). The effect of exposure on alpine treeline position: a case study from the High Sudetes, Czech Republic. *Arctic, Antarctic, and Alpine Research*, 40(4), 751-760.
- Treml, V., & Chuman, T. (2015). Ecotonal dynamics of the altitudinal forest limit are affected by terrain and vegetation structure variables: an example from the Sudetes Mountains in Central Europe. *Arctic, Antarctic, and Alpine Research*, 47(1), 133-146.
- Treml, V., & Migoń, P. (2015). Controlling factors limiting timberline position and shifts in the Sudetes: A review. *Geographia Polonica*, 88(2), 55-70.
- Treml, V., Banaš, M. 2000. Alpine Timberline in the High Sudetes. *Acta Universitatis Carolinae, Geographica*: 83-99. Karolinum, Praha.
- Treml, V., Šenfeldr, M., Chuman, T., Ponocná, T., & Demková, K. (2016). Twentieth century treeline ecotone advance in the Sudetes Mountains (Central Europe) was induced by agricultural land abandonment rather than climate change. *Journal of Vegetation Science*, 27(6), 1209-1221.
- Tyree, M.T., Sperry, J.S. 1989. Vulnerability of xylem to cavitation and metabolism. *Annual review of plant physiology and plant Molecular Biology*, 40: 19 – 38.