

Prach K.: O sukcesi a jejím využití v ekologii obnovy IV. Vše se mění (panta rhei)
(Živa 2026, 4: 199–202)

Literatura:

Dostálek J., Frantík T. (2011). Response of dry grassland vegetation to fluctuations in weather conditions: a 9-year case study in Prague (Czech Republic). *Biologia* 66/5: 837—847.

Fischer F.M., Chytrý K., Těšitel J., Danihelka J., Chytrý M. (2020). Weather fluctuations drive short-term dynamics and long-term stability in plant communities: A 25-year study in a Central European dry grassland. *J. Veg. Sci.* 31:711–721.

Holubičková B. (1959). Příspěvek ke studiu rašeliništní vegetace. I. Mokré louky u Třeboně. *Sborník Vysoké Školy Zemědělské v Praze* 1959:257–285

Hédl R. (2004). Vegetation of beech forests in the Rychlebské Mountains, Czech Republic, re-inspected after 60 years with assessment of environmental changes. *Plant Ecology* 170: 243–265.

Hroudová Z., Prach K. (1986). Vegetational changes on permanent plots in a steppe community. *Preslia* 58: 55–62.

Hroudová Z., Zákravský P. (1999). Vegetation dynamics in a fishpond littoral related to human impact. *Hydrobiologia* 415: 139–145.

Hroudová Z., Prach K., Zákravský P., Petřík P., Wild J. (2023). Jak reagují teplomilné travinné porosty v Českém krasu na suché roky? *Živa* 2/2023.

Chytrý M., Danihelka J. (1993). Long-term changes in the field layer of oak and oak-hornbeam forests under the impact of deer and mouflon. *Folia Geobotanica et Phytotaxonomica* 28: 225–245.

Klinkovská K. a kol. (2025). Half a Century of Temperate Non-Forest Vegetation Changes: No Net Loss in Species Richness, but Considerable Shifts in Taxonomic and Functional Composition. *Global Change Biology* 2025; 31:e70030.

Knollová I. a kol. (2024). ReSurveyEurope: A database of resurveyed vegetation plots in Europe. *J Veg Sci.* 2024; 35:e13235.

Lepš J., Lisner A. (2025). Resistance and Resilience of Species Composition: Thirty Years of Experimental Mismanagement and Subsequent Restoration in a Species Rich Meadow. *Ecology and Evolution*, 2025; 15:e70923.

Navrátilová J., Hájek M., Navrátil J., Hájková P., Frazier R.J. (2017). Convergence and impoverishment of fen communities in a eutrophicated agricultural landscape of the Czech Republic. *Applied Vegetation Science* 20 :225–235.

Pärtel M. a kol. (2025). Global impoverishment of natural vegetation revealed by dark diversity. *Nature* 641: 917–924.

- Prach J., Kopecký M. (2018). Landscape-scale vegetation homogenization in Central European sub-montane forests over the past 50 years. *Appl. Veg. Sci.* 21:373–384.
- Prach K. (2008): Vegetation changes in a wet meadow complex during the past half century. *Folia Geobotanica* 43: 119-130.
- Prach K., Vítovcová K., Řehouňková K., Královec J. (2021). Three decades of vegetation changes in a grassland after cessation of intensive fertilization. *Preslia* 93: 169–179.
- Prausová R., Doležal J., Rejmánek M. (2020). Nine decades of major compositional changes in a Central European beech forest protected area. *Plant Ecology* 221:1005–1016
- Skálová H., Hadincová V., Krahulec F., Pecháčková S., Herben T. (2022). Dynamics of a mountain grassland: Environment predicts long-term trends, while species' traits predict short-term fluctuations. *J. Veg. Sci.* 33:e13138.
- Tichý L. a kol. (2026). Successional changes of dry grasslands of rocky outcrops at Dyje/Thaya river valley (Czech Republic/Austria). *Preslia* 98: 169-185.
- Vild O. a kol. (2024). Long-term shift towards shady and nutrient-rich habitats in Central European temperate forests. *New Phytologist* 242: 1018–1028.